

Fourier Series:

Sines & Cosines:

Example:

$$f(x) = b_1 \sin \frac{px}{L} + b_2 \sin \frac{2px}{L} + \dots$$

Or:

$$f(x) = \sum_{n=1}^{\infty} b_n \sin \frac{npx}{L}$$

Now, a property of sines is that:

$$\int_{-L}^{+L} \sin \frac{npx}{L} \sin \frac{mpx}{L} dx = \begin{cases} 0 & m \neq n \\ L & m = n \end{cases}$$

On the interval $-L \leq x \leq +L$

Now, to find the coefficient b_1 , multiply both sides by $\sin \frac{px}{L}$, and integrate:

$$\int_{-L}^{+L} \sin \frac{px}{L} f(x) dx = \int_{-L}^{+L} b_1 \sin \frac{px}{L} \sin \frac{px}{L} dx + \int_{-L}^{+L} b_2 \sin \frac{2px}{L} \sin \frac{px}{L} dx + \dots$$

Which is all zero, except the first term, which is half the interval length. Hence:

$$b_1 = \frac{1}{L} \int_{-L}^{+L} \sin \frac{px}{L} f(x) dx$$

Hence, one should see, more generally:

$$b_n = \frac{1}{L} \int_{-L}^{+L} f(x) \sin \frac{npx}{L} dx \quad n = 1, 2, 3, \dots$$

Similarly for cosines:

If:

$$f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos \frac{npx}{L}$$

Then:

$$a_0 = \frac{1}{2L} \int_{-L}^{+L} f(x) dx$$
$$a_n = \frac{1}{L} \int_{-L}^{+L} f(x) \cos \frac{npx}{L} dx \quad n = 1, 2, 3, \dots$$

Technically, there is a b_0 term – but it is zero. Note that the a_0 isn't zero, and is divided by $2L$.

Now, for a general function:

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right) \quad \text{on } -L \leq x \leq +L$$

Where:

$$\begin{aligned} b_n &= \frac{1}{L} \int_{-L}^{+L} f(x) \sin \frac{n\pi x}{L} dx \\ a_0 &= \frac{1}{2L} \int_{-L}^{+L} f(x) dx \\ a_n &= \frac{1}{L} \int_{-L}^{+L} f(x) \cos \frac{n\pi x}{L} dx \end{aligned} \quad n = 1, 2, 3, \dots$$

Note that you divide a_n & b_n by half the length of the interval/integral.

For Fourier series to work, $f(x)$ must satisfy **Dirichlet's conditions**:

- $f(x)$ must be single valued & have finite number of discontinuities;
- $\int_{-L}^{+L} |f(x)| dx$ must be finite

Hence the Fourier series will *converge* to $f(x)$.

Fourier series use the property of the orthogonality between sines and cosines:

$$\int_{-L}^{+L} \cos \frac{n\pi x}{L} \sin \frac{m\pi x}{L} dx = 0$$

Complex Exponentials:

Two complex functions are orthogonal on $a \leq x \leq b$ if:

$$\int_a^b u^*(x)v(x) dx = 0$$

Hence, the two complex exponentials are orthogonal:

$$\int_{-L}^{+L} \left(e^{\frac{in\pi x}{L}} \right)^* \left(e^{\frac{im\pi x}{L}} \right) dx = \begin{cases} 0 & m \neq n \\ 2L & m = n \end{cases}$$

Thus, if:

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{\frac{in\pi x}{L}}$$

Then: for $-L \leq x \leq +L$

$$c_n = \frac{1}{2L} \int_{-L}^{+L} e^{-\frac{in\pi x}{L}} dx$$